

School Immunization Regulations

* **2017-18 School Immunization Requirements** - *Please note: the State of Pennsylvania has changed immunization requirements for school entry starting August 2017. The PA Department of Health has more information about those changes here:*

*<http://www.health.pa.gov/My%20Health/Immunizations/Pages/2017---2018-School-Immunization-Requirements.aspx#>.
ACHD will update its regulations accordingly to comply with state law.*

Key changes:

- (1) An additional dose of meningococcal vaccine for 12th graders*
- (2) A fourth dose of the polio vaccine on or after a child's fourth birthday (a fourth dose is not necessary if the third dose was administered at age four years or older and at least six months after the previous dose)*
- (3) A shortened provisional enrollment period (from 8 months to 5 days)*
- (4) A requirement that students not fully vaccinated by the end of the provisional enrollment period submit a plan from their physician stating when they will be up to date with vaccines.*

Immunizations Required For Students In Grades K-12

- ✓ 4 doses of **tetanus, diphtheria, and acellular pertussis** (1 dose must have been given on or after the 4th birthday)

Note: If series is started after 7 years of age, only 3 doses are required.

- ✓ ~~3 doses~~ 4 doses of **polio vaccine** (4th dose on or after 4th birthday and at least 6 months after previous dose given) **see above*

Note: If series is started after 7 years of age, only 3 doses are required.

- ✓ 2 doses of **measles, mumps, and rubella vaccine (MMR)**

- ✓ 3 doses of **hepatitis B vaccine**

- ✓ 2 doses of **varicella vaccine**, or written statement from physician/designee indicating month and year of disease or serologic proof of immunity

Additional Immunizations Required For Students In Grades 7-12

- ✓ 1 dose of **tetanus/diphtheria/pertussis vaccine (Tdap)**

- ✓ ~~1 dose~~ 2 doses of **meningitis**, one for entry into 7th grade and a second for entry into 12th grade. **see above*



Hartley Rose Building
425 First Avenue
Pittsburgh, PA 15219
412.578.8062
www.achd.net

Vaccinations for Preteens and Teens, Age 11–19 Years

Getting immunized is a lifelong, life-protecting job. Make sure you and your healthcare provider keep your immunizations up to date. Check to be sure you've had all the vaccinations you need.

Vaccine	Do you need it?
Chickenpox (varicella; Var)	If you haven't been vaccinated and haven't had chickenpox, you need 2 doses of this vaccine. Anybody who was vaccinated with only 1 dose should get a second dose.
Hepatitis A (HepA)	You need 2 doses of hepatitis A vaccine if you would like to be protected from this disease or if you have a risk factor (such as international travel) for hepatitis A. Check with your healthcare provider to find out if you need this vaccine.
Hepatitis B (HepB)	This vaccine is recommended for all people age 0–18 years. You need a hepatitis B vaccine series if you have not already received it.
Human papillomavirus (HPV)	All preteens and teens age 11 and older need a series of doses of HPV vaccine. The vaccine protects against HPV, the most common cause of cervical cancer. It also protects against some other types of cancers, such as cancer of the anus, penis, and throat. HPV vaccine also protects against genital warts.
Influenza (Flu)	Everyone age 6 months and older needs annual influenza vaccination every fall or winter and for the rest of their lives.
Measles, mumps, rubella (MMR)	You need 2 doses of MMR vaccine if you have not already received them. MMR vaccine is usually given in childhood.
Meningococcal (MenACWY [MCV4], MenB)	All preteens and teens need 2 doses of MenACWY vaccine, the first at age 11–12 years and the second at age 16 years. Teens can also receive a MenB vaccine series at ages 16–18 years. If you are a first-year college student living in a residence hall, you need a dose of MenACWY if you never received it or received it when you were younger than 16. Check with your healthcare provider.
Pneumococcal (Pneumovax [polysaccharide vaccine, PPSV])	Do you have an ongoing health condition? If so, check with your healthcare provider to find out if you need one or both of the pneumococcal vaccines.
Polio (IPV)	You need a series of at least 3 doses of polio vaccine if you have not already received them. Polio vaccine is usually given in childhood.
Tetanus, diphtheria, and whooping cough (pertussis; Tdap)	All preteens and teens (and adults!) need a dose of Tdap vaccine, a vaccine that protects you from tetanus, diphtheria, and whooping cough (pertussis). After getting a dose of Tdap, you will need a tetanus-diphtheria (Td) shot every ten years. If you become pregnant, however, you will need another dose of Tdap during the pregnancy, preferably during the third trimester.

Will you be traveling outside the United States? Visit the Centers for Disease Control and Prevention's (CDC) website at wwwnc.cdc.gov/travel/destinations/list for travel information, or consult a travel clinic.