

Allies For Health + Wellbeing

5913 Penn Avenue, Pittsburgh, PA 15206 Main: 412.345.7456 Clinic: 412.248.0550

Central Outreach Wellness Center

Timber Court Building 127 Anderson St., Ste. 10, Pittsburgh, PA 15212 412.515.0000

Persad Center

5301 Butler St #100, Pittsburgh, PA 15201 412.441.9786

Planned Parenthood of Western PA

933 Liberty Avenue, Pittsburgh, PA 15222 412.434.8971

Prevention Point 460 Melwood Ave #205, Pittsburgh, PA 15213 412-247-3404

SisTers PHG 2014 Monongahela Ave, Pittsburgh, PA 15218 412-259-3091

Trans Buddy PGH 412.944.4261 transbuddypgh@gmail.com



Allegheny County Public Health Clinic

1908 Wylie Avenue (Hill District) Pittsburgh, PA 15213

412-578-8081

www.achd.net

Clinic Hours

Monday, Tuesday, Thursday, Friday 8:30 a.m. -3:30 p.m. Wednesday 12:30 p.m. -7:30 p.m.

Registration starts 30 minutes earlier



Transgender Health and Safer Sex



SAFER SEX TIPS

KNOW YOUR RISK

Your risk of getting an STD is determined by the types of sex you have. For example, anal sex without a condom can put you at higher risk of getting an STD because the anus does not selflubricate. You can also get many of the same STDs through oral sex (like chlamydia, gonorrhea, syphilis, and herpes) that you can get through anal or vaginal sex.

USE CONDOMS AND LUBRICANT

By reducing friction during sex, lubricant can help reduce the risk of small, invisible tears happening in the rectum or vagina. Lubricant is especially important for anal sex, or if you or your partner experiences vaginal dryness. There are many types of lubricant, but remember that only water-based lubricant can be used with latex condoms.

ASK ABOUT PREP

PrEP (pre-exposure prophylaxis) is one pill a day that can reduce your chance of getting HIV. When taken as directed, PrEP can reduce the risk of getting HIV from sex by more than 90%, and for those who inject drugs by more than 70%. Your risk of getting HIV from sex can be even lower if you use condoms as the same time as PrEP.

MEDICAL CARE

Many transgender people have difficulty accessing healthcare for a number of reasons,. Here are some tips that can help:

- 1. Find a provider that respects your identity and offers the treatment that you need. Listed on the back of this pamphlet are several medical resources that specialize in transgender healthcare.
- 2. Be honest with your provider about the type of sex you are having, and how many partners you are having sex with. This will help your provider determine your risk for STDs, and help you find the best way to prevent them.
- 3.Know that the type of screening tests you need will depend on the anatomy you have.
- 4. Bring a buddy. If it makes you feel safer or more comfortable, bring a trusted friend to your appointment. You can also request a trained advocate from Trans Buddy PGH (contact information listed on back) to attend your appointment with you at no cost.

HIV AND HEPATITIS C

- Transgender people may use needles for a variety of reasons, including silicone or hormone injection. If you share needles with others, it is possible to get STDs like HIV or hepatitis C.
- HIV is a virus that attacks your immune system that cannot be cured, and will lead to severe illness and death if left untreated.
- Hepatitis C is a virus that attacks the liver, causing liver damage.
- You should always use a new or unused needle when possible.
- When you are done with the needle, you can put it in a hard plastic container with a lid, such as a laundry detergent bottle. Once it is full, tape the lid on tightly. You can throw it in the regular garbage.
- You can also get a needle container from some needle exchange programs, like **Prevention Point** (contact information is listed on "Resources" section).